

The R 'sampling' package

Alina Matei and Yves Tillé
University of Neuchâtel

Cardiff, Q2006
April 2006

The R language

- ▶ Shareware available on <http://cran.r-project.org/>

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network
- ▶ Installation: 10 minutes

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network
- ▶ Installation: 10 minutes
- ▶ All the manuels are available in pdf

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network
- ▶ Installation: 10 minutes
- ▶ All the manuels are available in pdf
- ▶ Everyone can write an additional package (600 packages are available)

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network
- ▶ Installation: 10 minutes
- ▶ All the manuels are available in pdf
- ▶ Everyone can write an additional package (600 packages are available)
- ▶ Packages are loaded directly from R.

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network
- ▶ Installation: 10 minutes
- ▶ All the manuels are available in pdf
- ▶ Everyone can write an additional package (600 packages are available)
- ▶ Packages are loaded directly from R.
- ▶ The manual of the package is available online and in pdf.

The R language

- ▶ Shareware available on <http://cran.r-project.org/>
- ▶ The Comprehensive R Archive Network
- ▶ Installation: 10 minutes
- ▶ All the manuels are available in pdf
- ▶ Everyone can write an additional package (600 packages are available)
- ▶ Packages are loaded directly from R.
- ▶ The manual of the package is available online and in pdf.
- ▶ Package 'sampling' written by Matei and Tillé.

Continuous distributions

- ▶ EFTA course for public statisticians (April 2005).

Continuous distributions

- ▶ EFTA course for public statisticians (April 2005).
- ▶ Objective : to apply directly the theory with the R language.

Continuous distributions

- ▶ EFTA course for public statisticians (April 2005).
- ▶ Objective : to apply directly the theory with the R language.
- ▶ Theory + Exercices with a laptop and R.

Continuous distributions

- ▶ EFTA course for public statisticians (April 2005).
- ▶ Objective : to apply directly the theory with the R language.
- ▶ Theory + Exercices with a laptop and R.
- ▶ Writing of a large set of procedures.

Continuous distributions

- ▶ EFTA course for public statisticians (April 2005).
- ▶ Objective : to apply directly the theory with the R language.
- ▶ Theory + Exercices with a laptop and R.
- ▶ Writing of a large set of procedures.
- ▶ Finally, decision of submitting the package to the CRAN.

Content of the package

- ▶ Stratification, two-stage, unequal probabilities, balanced sampling

Content of the package

- ▶ Stratification, two-stage, unequal probabilities, balanced sampling
- ▶ Estimation: calibration and regression estimator

Content of the package

- ▶ Stratification, two-stage, unequal probabilities, balanced sampling
- ▶ Estimation: calibration and regression estimator
- ▶ Tools : computation of inclusion probabilities, crossing strata

Content of the package

- ▶ Stratification, two-stage, unequal probabilities, balanced sampling
- ▶ Estimation: calibration and regression estimator
- ▶ Tools : computation of inclusion probabilities, crossing strata
- ▶ Data bases, Swiss municipalities, Belgian municipalities.

Tools

- ▶ `writesample`: return the list of all the samples of fixed sample size

Tools

- ▶ `writesample`: return the list of all the samples of fixed sample size
- ▶ `cleanstrata`: renumbering of the strata

Tools

- ▶ `writesample`: return the list of all the samples of fixed sample size
- ▶ `cleanstrata`: renumbering of the strata
- ▶ `disjonctive` return a matrix with 0 and 1 that is the disjonctive representation of the stratum.

Tools

- ▶ `writesample`: return the list of all the samples of fixed sample size
- ▶ `cleanstrata`: renumbering of the strata
- ▶ `disjonctive` return a matrix with 0 and 1 that is the disjunctive representation of the stratum.
- ▶ `inclusionprobabilities`: compute unequal inclusion probabilities from an auxiliary variable variable.

Data bases

- ▶ MU284 A data frame with 284 municipalities on the following 11 variables : populations, political results.

Data bases

- ▶ MU284 A data frame with 284 municipalities on the following 11 variables : populations, political results.
- ▶ `swissmunicipalities`: 2896 Swiss municipalities. Surfaces and population.

Data bases

- ▶ MU284 A data frame with 284 municipalities on the following 11 variables : populations, political results.
- ▶ `swissmunicipalities`: 2896 Swiss municipalities. Surfaces and population.
- ▶ `belgianmunicipalities`: 589 Belgian municipalities 11 variables, population and taxes.

Simple random sampling

- ▶ `srswor`: Simple random sampling with replacement.

Simple random sampling

- ▶ `srswor`: Simple random sampling with replacement.
- ▶ `srswor1`: Simple random sampling without replacement (sequential method).

Simple random sampling

- ▶ `srswor`: Simple random sampling with replacement.
- ▶ `srswor1`: Simple random sampling without replacement (sequential method).
- ▶ `srswr`: Simple random sampling with replacement.

Unequal probability sampling

- ▶ UPbrewer,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,
- ▶ UPmultinomial,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,
- ▶ UPmultinomial,
- ▶ UPpivotal, UPrandompivotal,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,
- ▶ UPmultinomial,
- ▶ UPpivotal, UPrandompivotal,
- ▶ UPpoisson,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,
- ▶ UPmultinomial,
- ▶ UPpivotal, UPrandompivotal,
- ▶ UPpoisson,
- ▶ UPsampford,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,
- ▶ UPmultinomial,
- ▶ UPpivotal, UPrandompivotal,
- ▶ UPpoisson,
- ▶ UPsampford,
- ▶ UPsystematic, UPrandomsystematic, UPsystematicpi2,

Unequal probability sampling

- ▶ UPbrewer,
- ▶ UPmaxentropy, (set of function)
- ▶ UPmidzuno, UPmidzunopi2,
- ▶ UPmultinomial,
- ▶ UPpivotal, UPrandompivotal,
- ▶ UPpoisson,
- ▶ UPsampford,
- ▶ UPsystematic, UPrandomsystematic, UPsystematicpi2,
- ▶ UPtille, UPtillepi2,

Balanced sampling

- ▶ Design that satisfies the balancing equations

$$\sum_{k \in S} \frac{\mathbf{x}_k}{\pi_k} = \sum_{k \in U} \mathbf{x}_k,$$

where \mathbf{x}_k is a vector of auxiliary variables.

Balanced sampling

- ▶ Design that satisfies the balancing equations

$$\sum_{k \in S} \frac{\mathbf{x}_k}{\pi_k} = \sum_{k \in U} \mathbf{x}_k,$$

where \mathbf{x}_k is a vector of auxiliary variables.

- ▶ Cube algorithm: flight phase and landing phase.

Balanced sampling

- ▶ Design that satisfies the balancing equations

$$\sum_{k \in S} \frac{\mathbf{x}_k}{\pi_k} = \sum_{k \in U} \mathbf{x}_k,$$

where \mathbf{x}_k is a vector of auxiliary variables.

- ▶ Cube algorithm: flight phase and landing phase.
- ▶ samplecube, fastflightcube, landingcube

Balanced sampling

- ▶ Design that satisfies the balancing equations

$$\sum_{k \in S} \frac{\mathbf{x}_k}{\pi_k} = \sum_{k \in U} \mathbf{x}_k,$$

where \mathbf{x}_k is a vector of auxiliary variables.

- ▶ Cube algorithm: flight phase and landing phase.
- ▶ `samplecube`, `fastflightcube`, `landingcube`
- ▶ Complex survey `balancedstratification`
`balancedcluster` `balancedtwostage`